

An Integrated Analysis of Maternal-Infant Sleep, Breastfeeding, and Sudden Infant Death Syndrome Research Supporting a Balanced Discourse

Journal of Human Lactation

1–11

© The Author(s) 2019

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/0890334419851797

journals.sagepub.com/home/jhl



Kathleen A. Marinelli, MD, IBCLC, FABM¹ ,
Helen L. Ball, BSc (Hons), MA, PhD²,
James J. McKenna, MA, PhD³,
and Peter S. Blair, BSc (Hons), MSc, PhD⁴

Abstract

Breastfeeding and the place of sleep for the mother and the infant have been controversial internationally due to reported concerns regarding infant deaths despite the known benefits of exclusive and prolonged breastfeeding, which are increased by breastfeeding at night. The aims of this integrated analysis were to (a) review breastfeeding and maternal and infant sleep research literature via historical, epidemiological, anthropological, and methodological lenses; (b) use this information to determine where we are currently in safeguarding both infant lives and breastfeeding; and (c) postulate the direction that research might take from this point forward to improve our knowledge and inform our policy and practice. Despite well-meaning but unsuccessful campaigns in some countries to dissuade parents from sleeping with their babies, many breastfeeding mothers and caregivers do sleep with their infants whether intentionally or unintentionally. Taking cultural contexts and socio-ecological circumstances into consideration, data supports policies to counsel parents and caregivers on safe sleep practices, including bed sharing in non-hazardous circumstances, particularly in the absence of parental smoking, recent parental alcohol consumption, or sleeping next to an adult on a sofa. Further research with appropriate methodology is needed to drill down on actual rates of infant deaths, paying close attention to the definitions of deaths, the circumstances of the deaths, and confounding factors, in order to ensure we have the best information with which to derive public health policy. Introduction and use of the concept of “breastsleeping” is a plausible way to remove the negative connotations of “co-sleeping” and redirect ongoing data-driven discussions and education of best practices of breastfeeding and sleep.

Keywords

bed-sharing, breastfeeding, co-sleeping, epidemiological methods, infant behavior, lactation, maternal behavior, mother-infant dyad

Background

The matter of breastfeeding and the place of sleep of the mother and the infant has been and remains a contentious issue in maternal-child health globally. The aims of this paper were to review the literature associated with this topic using an historical, epidemiological, anthropological, and methodological lens and then, with the information determine where we are now in safeguarding both infant lives and breastfeeding, as well as the direction that research might take from this point forward to improve knowledge and inform our policy and practice.

¹University of Connecticut School of Medicine, Farmington, CT, USA

²Infancy & Sleep Centre, Department of Anthropology, Durham University, Durham, UK

³Mother-Baby Behavioral Sleep Laboratory, Department of Anthropology, University of Notre Dame, Notre Dame, Indiana, USA

⁴Centre of Academic Child Health, Bristol Medical School, University of Bristol, Bristol, UK

Date received: 30 April 2019; accepted: 30 April 2019

Corresponding Author:

Kathleen A. Marinelli MD, IBCLC, FABM, Clinical Professor of Pediatrics, University of Connecticut School of Medicine, Farmington CT, USA, Connecticut Children's Medical Center Division of Neonatology, Hartford CT, USA.

Email: kathleen.marinelli@cox.net

History and Background to Current Debate

This issue is of relevance for mothers and their babies and other caregivers throughout the world. The bed-sharing debate ignited in the late twentieth century during a period of intense research that sought explanations for unexplained infant deaths (known as Sudden Infant Death Syndrome [SIDS]) that occurred during sleep and peaked during the second to third month of life. Although the cause of SIDS remains unknown, epidemiological studies identified a strong association with infant sleep position, leading to campaigns that informed parents to put babies to sleep on their backs (Gilbert, Salanti, Harden, & See, 2005). Other infant care practices in the sleep environment were intensely scrutinized in Western countries, for example the use of pacifiers, soft bedding, and infant head coverings.

Anthropologists noted how poorly Western sleep arrangements met the human infant's unique biological and behavioral needs (McKenna, 1986; Konner & Super, 1987), with babies predominantly sleeping alone in a crib in a separate room from their parents. As placental mammals, humans produce helpless young who require prolonged maternal post-natal care and lactation. This is especially characteristic of the human infant who is born neurologically the least mature primate of, the most dependent on the caregiver for bodily regulation, and for the longest period of time. Human milk follows the typical primate composition: low in protein and fat, but high in sugar (Jelliffe & Jelliffe, 1978). Cross-cultural studies emphasize that in most traditional societies infants are maintained in physical contact with their mother day and night, experience frequent arousals during sleep, and suckle on demand throughout the first year of life (Ball, 2007). In contrast, social and cultural changes in industrial and post-industrial societies have encouraged solitary and prolonged sleep periods from an early postnatal age, which are now considered characteristic of Western cultures (McKenna, Ball, & Gettler, 2007).

Against this backdrop, anthropologists hypothesized that one explanation for SIDS could be the unusual practice of separating babies and mothers at night (McKenna, 1986; McKenna et al., 1993; Konner & Super, 1987). That infants undergo dramatic changes in their breathing control at around 3 months of age makes them particularly vulnerable to unpredictable breathing cessation (Mosko, Richard, & McKenna, 1997a). SIDS researchers were investigating suppressed infant arousals and breathing pauses (apnea), as potential precursors to unexpected infant deaths when McKenna et al. (1993) proposed that infants experiencing close sleep contact may be protected from apneic pauses and blunted arousal from sleep by maternal sounds, movements, and breathing. In a series of polysomnographic studies of breastfeeding mothers and babies, McKenna and colleagues demonstrated that sleep contact between them promoted regular night-time interaction and lighter stages of sleep, with

Key Messages

- Sudden Infant Death Syndrome studies have often been interpreted as consistent with policies to exclude breastfeeding mothers from sleeping with their infants, even in the absence of risk factors for infant death.
- Except for policy in the United Kingdom, the primary source of information to guide families on safe sleeping and breastfeeding that does not separate the mother and the child comes from breastfeeding support organizations.
- In the context of “breastsleeping” we can emphasize the magnitude of risks surrounding unsafe sleeping practices involving alcohol, drugs, and sofas or chairs, and call for a more coordinated approach with public health strategists on how best to care for infants and keep them safe.

fewer obstructive apneas (McKenna, Mosko, Dungy, & McAninch, 1990; Mosko, McKenna, Dickel, & Hunt, 1993; Mosko, Richard, & McKenna, 1997a; 1997b). This led them to propose that mother-baby co-sleeping was adaptive for breastfeeding dyads and helped to protect babies from SIDS. However, epidemiologists studying sudden infant deaths were encountering different patterns in their data that indicated mother-baby co-sleeping was associated with increased risk (Mitchell & Scragg, 1993). To them, the notion that mothers and babies might choose to sleep together was inherently problematic. The ensuing discourse, debate, and disagreements about the role of mother-infant sleep contact in reducing or increasing sudden and unexpected infant deaths has continued for over 3 decades.

The progression of the discussion over co-sleeping, breastfeeding, and SIDS has been an iterative one involving negotiation and re-negotiation of research foci, data collection methods, variable definitions, and the ever-closer interrogation of more detailed data sets. The more carefully the details are examined, the clearer it becomes that the answer to the simple question of whether mother-infant sleep contact is a good or bad thing is “it depends” (Ball, 2017a).

Epidemiology of Bed Sharing

Prolonged physical contact between parents and infants during sleep is normal infant care behavior in many different cultures and, despite previous assumptions to the contrary, is commonly practiced in Western societies. In England, almost half of all neonates bed-share at some time with their parents, and a fifth of infants are brought into the parental bed on a regular basis over the first year of life (Blair & Ball, 2004). In the US, bed-sharing prevalence peaks in early infancy and declines with increasing age (Blair & Ball, 2004; Blair, Heron, & Fleming, 2010; Colson et al., 2013; McCoy et al.,

2004). Older data from the US National Infant Sleep Position Study (Willinger, Ko, & Hoffman, 2003) found that over 40% of 8000 parents reported that their infants slept with them in an adult bed. The strongest predictors of bed-sharing in the US were being black or Asian, breastfeeding, having a mother younger than 18 years old, and low household income. An increase in bed-sharing between 1993 and 2000 was associated with maternal age >18 years; self-identification as white or Asian; infant age older than 8 weeks; and term infants with normal birth weight. More recently in the United States, about 21% of all mothers and more than 25% of Hispanic mothers, reported bed sharing for some or all of the night (Smith et al., 2016). This is likely to represent a significant undercount, especially in the United States, given the fact that the condemnation of bed sharing and bed-sharing families there makes it difficult for families to be honest about where their infants actually sleep.

Bed Sharing and Breastfeeding

A strong relationship between bed sharing and breastfeeding has been demonstrated in multiple studies, suggesting that a “Never Bed-share” message may not only hinder maternal and child health promotion by impeding breastfeeding (Ball, 2003; Bartick & Smith, 2014), but also SIDS reduction itself, as breastfeeding has been associated with a greatly reduced risk of SIDS (Hauck, Thompson, Tanabe, Moon, & Vennemann, 2011; Thompson et al., 2017). It has also become clear that although bed-sharing has been associated with particular cultural contexts and socio-ecological circumstances (Luijk et al., 2013; Salm Ward & Doering, 2014) SIDS rates are extremely low in some of these settings (e.g., Ball et al., 2009).

Epidemiology of Sudden Infant Death Syndrome and Bed Sharing

Although flagged as a potential risk factor 3 decades ago (Mitchell & Scragg, 1993), the definitions of “bed-sharing deaths” varied widely, encompassing sofa sharing, sleep sharing with siblings or pets, and babies returned to a crib (Côté, 2006). Furthermore, control families’ reports of “bed-sharing” were not collected using standard definitions, and ranged from “usual behavior,” to sleep location on a particular night (or part of the night), to ever sharing a sleep surface. Data were not comparable between studies, or even between cases and controls in the same study (Ball, 2007; Ball, Hooker, & Kelly, 1999). Potential interactions between bed-sharing and hazardous circumstances, such as infants sleeping next to parents who smoked, drank alcohol, or used a sofa for the sleep started to emerge (Blair et al., 2009), but carried different weight in national risk reduction campaigns. The relationship between bed sharing and SIDS was revealed to be more complex than initially assumed (Ball &

Volpe, 2013; Fetherston & Leach, 2012) and the approach on how to advise parents diverged into both strict guidance to avoid bed sharing and acknowledging that bed sharing happens and discussing when this may or may not be appropriate.

Although commonly known as “cot death” or “crib death,” SIDS can occur in any infant sleeping environment and has increasingly been discovered to occur in shared sleeping spaces more often than expected. Recent observational case-control studies suggest that as many as half of SIDS deaths occur when infants sleep alongside an adult (Vennemann et al., 2012). This rather alarming proportional rise in SIDS deaths outside the cot has led some countries to advise against bed-sharing, including, the American Academy of Pediatrics (AAP) since 2015 (American Academy of Pediatrics, 2005; Moon, 2016a). A meta-analysis of 11 SIDS case-control studies published in 2012 showed a pooled 3-fold increased risk associated with bed-sharing, although this did not reach significance in older infants (> 12 weeks) or those not exposed to tobacco smoke (Vennemann et al., 2012). Longitudinal data from Avon, UK of 300 consecutive SIDS deaths over a 20-year period show that the proportional rise in bed-sharing SIDS deaths does not equate to a numerical increase (Blair, Sidebotham, Berry, Evans, & Fleming, 2006). The striking feature in this unique dataset is the 7-fold fall in deaths occurring in the cot (Figure 1). SIDS deaths in the parental bed also fell by half over this time period, but increased proportionally as part of the whole. Why the “Back to Sleep” campaign (American Academy of Pediatrics, 2019) was less effective among bed-sharing deaths is not clear, although data from the Avon cohort (Blair et al., 2006) and subsequent studies (Blair et al., 2014) suggested that placing infants prone to sleep was far more common among infants sleeping alone rather than those sleeping with someone. This may partly explain the inherent protection of breastfeeding against SIDS, in that to initiate or enable this process infants are more likely to be placed supine, and that the exposure risk while bed-sharing may thus lie elsewhere. Notably, the only sleeping environment in which SIDS increased during these 20 years was infants sleeping next to a parent on a sofa. This is particularly important to observe because any caregiver seeking to avoid the potential risk of bringing the baby into bed to breastfeed and inadvertently falling asleep may put the baby at greater risk by getting out of bed and sitting in a chair or on a sofa.

Significant Interactions Providing Hazardous Exposure to the Infant

The interaction between maternal smoking and bed-sharing as a risk for SIDS was first identified by Mitchell and colleagues’ large New Zealand study (Scragg et al., 1993) and has been confirmed in their most recent study (Mitchell




Figure 1. SIDS deaths by sleeping environment (300 consecutive SIDS deaths in Avon, UK between 1984 and 2003).

et al., 2017). The risk among infants bed-sharing next to mothers who smoked was more than 4-fold ($OR = 4.55$; 95% CI [2.63–7.88]) compared to no risk among infants sleeping next to nonsmoking mothers ($OR = 0.98$; 95% CI [0.44–2.18]). Similar findings were observed in subsequent studies. It is not clear why this exposure would put the infant at risk; innate vulnerability due to fetal exposure to tobacco smoke during pregnancy postnatal prolonged passive exposure, or a proxy marker for some other unmeasured risk-taking parental behavior have all been postulated but little further evidence has been provided. Significant interactions have also been observed between bed-sharing and parental use of alcohol or drugs prior to the last sleep, and using a sofa to sleep with the infant (Blair et al., 2009). A combined analysis from two English studies (Blair et al., 2014) suggested an 18-fold increase in SIDS deaths if an infant sleeps next to an adult who drinks more than two units of alcohol or if an infant sleeps with an adult on a sofa. Both hazardous circumstances are suggestive that overlaying is a potential causal explanation for these SIDS deaths although it is difficult to verify a causal classification using current post-mortem techniques. Noticeably, the exposure of these two hazardous circumstances were very rare amongst the controls (< 1%) suggesting these practices are potentially lethal. Researchers report an observational study of nearly 8,000 sleep-related infant deaths in 24 US states between 2004–2012 found that over 800 occurred while the infants slept with an adult on a sofa (Rechtman, Colvin, Blair, & Moon, 2014). The diagnoses of these deaths were fairly evenly split between SIDS, ill-defined and accidental suffocation, and strangulation in bed (ASSB), suggesting that the prevalence of sofa-sharing deaths is far higher than first reported in observational SIDS

studies. Bed-sharing SIDS victims are younger than those infants found in cots/cribs, and other potential characteristics that may lead to increased risk include the use of pillows near the infant, parental exhaustion, vulnerable low birth-weight or premature infants, not breastfeeding, and lack of provision for a cot/crib, although further evidence is needed to confirm these factors.

The Risk of Bed-sharing in Non-Hazardous Circumstances

The combined analysis from England (Blair et al., 2014) suggested there was no risk of bed-sharing in the absence of three particular hazards: parental smoking; recent parental alcohol consumption; or sleeping next to an adult on the sofa. A subgroup analysis in Blair's study, limited to younger infants (under 12 weeks), increased the observed risk ($OR = 1.62$, 95% CI [0.96–2.73]) but this did not become significant. In contrast, a similar combined analysis showed a 5-fold increased risk associated with younger infants bed sharing in non-hazardous circumstances. However, the idealized reference group used was breastfed infants placed on their backs to sleep in a separate room by non-smoking parents, and in the absence of any other risk factors. This renders this quantification as uninterpretable as these two groups are not comparable despite this assumption (Carpenter et al., 2013). The AAP, in their review of the evidence to support their 2016 guidelines (Moon, 2016b), concluded that the data from these two different analyses do not support a definitive conclusion that bed sharing among the youngest infants is safe, even under less hazardous circumstances. In contrast a 2014 review of these two analyses by the independent National Institute for

Health and Care Excellence (NICE) (National Institute for Health and Care Excellence, 2014a) in the UK concluded that bed-sharing in itself is not causal and that parents need to be informed of the specific hazards associated with this practice.

Benefits of Bed Sharing

When parents are interviewed about sleeping with their baby they give various reasons for doing so (Ateah & Hamelin, 2008; Ball, 2002; Crane & Ball, 2016; Culver, 2009; McKenna & Volpe, 2007). Their answers express deeply-rooted cultural or religious beliefs and parenting philosophies, invoke the physiological links between lactation and night-time breastfeeding, and reflect the biological compulsion that drives the urge for close contact (Salm-Ward, 2015). On a practical level they explain that sleeping with the baby makes night-time care easier, and helps them to monitor the baby, provide comfort, and yet obtain sleep (Ball, 2002, 2003; Rudzik & Ball, 2016). Sometimes parents report having nowhere else to put their baby at night, or that they have fallen asleep with their baby unintentionally.

For breastfeeding mothers all the above reasons may apply, so it is unsurprising that the largest group of bed-sharers around the globe are breastfeeding mothers. Of 34 studies exploring maternal reasons for co-sleeping 26 reported breastfeeding as the key reason (Salm-Ward, 2015). Bed sharing facilitates night-time breastfeeding, and is associated with more frequent night-time feeds (which promotes milk production, taking advantage of the physiological prolactin surge), and with more months of breastfeeding. The observed association between bed-sharing and greater breastfeeding duration was demonstrated in Brazil, where researchers investigated breastfeeding outcomes at 12 months by interviewing mothers of infants at birth ($N = 4231$), 3 months, and 12 months about their feeding and sleeping arrangements. Breastfeeding prevalence at 12 months was 59% for those who bed shared (habitually for all or part of the night) at 3 months, and 44% for those who did not. Among infants exclusively breastfed at 3 months, 75% of bed-sharers were still breastfed at 12 months, versus 52% of non-sharers (Santos, Mota, Matijasevich, Barros, & Barros, 2009). Although the authors accepted these results as evidence that bed sharing protects against early weaning, the association did not reveal the direction of causality.

In the UK a similar association was found, with mothers being twice as likely to still be breastfeeding 6 months post-birth if they commenced bed sharing within the first 3 months, than if they did not; however, in this study, strength of intent to breastfeed had been assessed in early pregnancy and it was found that mothers who chose to bed share were those with the strongest intent to breastfeed to 6 months or beyond (Ball et al., 2016). These data are consistent with the interpretation that bed sharing is a strategy used by breastfeeding mothers to

reduce the costs (e.g., sleep disruption) of prolonged breastfeeding (Tully & Ball, 2013). Prevention of bed sharing is therefore likely to undermine breastfeeding goals (Ball, 2003).

With regard to maternal sleep, numerous studies have confirmed that although bed-sharing breastfeeding mothers wake frequently to feed, they also wake for shorter periods, fall back to sleep more rapidly (Mosko et al., 1997b), and achieve longer sleep duration (Quillin, & Glenn, 2004), when compared to those not bed sharing. Mothers choose to bed share to make night-time care easier and reduce sleep disruption, particularly when breastfeeding (Ateah et al., 2008; Rudzik et al., 2016). Although many breastfeeding mothers report having been told that bed sharing is “wrong,” almost every breastfeeding mother sometimes falls asleep with her baby, in bed, in a chair, or on a couch, regardless of whether or not she considers herself to be a “bed-sharer”. It is vital, therefore, that all health professionals who support breastfeeding mothers are well informed about the issues surrounding sleep sharing and can help new mothers make sense of how the research evidence relates to their own situations.

The Concept of Breastsleeping

In 2016 Mobbs and colleagues described the need for, and benefits of, immediate and sustained contact, including co-sleeping, to establish an appropriate foundation for optimal human infant breastfeeding, neonatal attachment, and brain growth. In further support of this model, and building upon existing knowledge of breastfeeding and sleep data, McKenna and Gettler (2016) proposed the concept of *breastsleeping*, with which they aimed to help resolve the bed-sharing debate and to distinguish between the known and potential differences between breastfeeding, bed-sharing dyads and non-breastfeeding, bed-sharing situations, particularly when breastfeeding and bed sharing is practiced in the absence of all known hazardous factors. They argued that, because breastfeeding “is so physiologically and behaviorally entwined and functionally interdependent with forms of co-sleeping” (McKenna & Gettler, 2016, p. 17) the term *breastsleeping* would encourage acknowledgment of the following components:

- (i) the critical role that immediate and sustained maternal contact plays in helping to establish optimal breastfeeding (Ball, 2008; 2017b; Ball & Russell, 2012);
- (ii) normal, human (species wide) infant sleep parameters should only be derived from studies of breast-sleeping dyads because of the ways maternal–infant contact affects the delivery of the mother’s milk, the milk’s ingestion, the infant’s concomitant and subsequent metabolism, and other physiological processes, maternal and infant sleep architecture, including arousal patterns (McKenna et al., 1990; Mosko et al.,

Table 1. American Academy of Pediatrics Guidance on Bed-sharing and Sudden Infant Death Syndrome and Other Sleep-Related Infant Deaths for the United States (2016).

It is recommended that infants sleep in the parents' room, close to the parents' bed but on a separate surface designed for infants, ideally for the 1st year of life, but at least for the first 6 months.

There is evidence that sleeping in the parents' room but on a separate surface decreases the risk of sudden infant death syndrome by as much as 50%. In addition, this arrangement is most likely to prevent suffocation, strangulation, and entrapment that may occur when the infant is sleeping in the adult bed.

Infants who are brought into the bed for feeding or comforting should be returned to their own crib or bassinet when the parent is ready to return to sleep.

Couches and armchairs are extremely dangerous places for infants. Sleeping on couches and armchairs places infants at extraordinarily high risk of infant death, including SIDS, suffocation through entrapment or wedging between seat cushions, or overlay if another person is also sharing this surface. Therefore, parents and other caregivers should be especially vigilant as to their wakefulness when feeding infants or lying with infants on these surfaces. Infants should never be placed on a couch or armchair for sleep.

The safest place for an infant to sleep is on a separate sleep surface designed for infants close to the parents' bed. However, the American Academy of Pediatrics acknowledges that parents frequently fall asleep while feeding the infant. Evidence suggests that it is less hazardous to fall asleep with the infant in the adult bed than on a sofa or armchair, should the parent fall asleep. It is important to note that a large percentage of infants who die of SIDS are found with their head covered by bedding. Therefore, no pillows, sheets, blankets, or any other items that could obstruct infant breathing or cause overheating should be in the bed. Parents should also follow safe sleep recommendations outlined elsewhere in this statement. Because there is evidence that the risk of bed-sharing is higher with longer duration, if the parent falls asleep while feeding the infant in bed, the infant should be placed on a separate sleep surface as soon as the parent awakens.

There are specific circumstances that, in case-control studies and case series, have been shown to substantially increase the risk of SIDS or unintentional injury or death while bed sharing, and these should be avoided at all times:

- Bed sharing with a term, normal-weight infant younger than 4 months, or an infant born preterm and/or with low birth weight, regardless of parental smoking status. Even for breastfed infants, there is an increased risk of SIDS when bed-sharing if younger than 4 months. This appears to be a particularly vulnerable time, so if parents choose to feed their infants younger than 4 months in bed, they should be especially vigilant to not fall asleep.
- Bed sharing with a current smoker (even if he or she does not smoke in bed) or if the mother smoked during pregnancy.
- Bed sharing with someone who is impaired in his or her alertness or ability to arouse because of fatigue or use of sedating medications (e.g., certain antidepressants, pain medications) or substances (e.g., alcohol, illicit drugs).
- Bed sharing with anyone who is not the infant's parent, including non-parental caregivers and other children.
- Bed sharing on a soft surface, such as a waterbed, old mattress, sofa, couch, or armchair.
- Bed sharing with soft bedding accessories, such as pillows or blankets.

Note. Adapted from "SIDS and Other Sleep-Related Infant Deaths: Updated 2016 Recommendations for a Safe Infant Sleeping Environment," by R. Y. Moon, & Task Force on Sudden Infant Death Syndrome, 2016, *Pediatrics*, 138(5), e20162938. Copyright 2016 by the American Academy of Pediatrics.

1997a, 1997b) as well as breastfeeding frequency and prolongation (Ball et al., 2006; 2016); and

- (iii) that breastsleeping by mother–infant pairs appears to involve substantially different behavioral and physiological characteristics compared with never-breastfeeding mothers and infants. This dyadic context must be more closely researched and more carefully understood, and potentially distinguished as a separate epidemiological category for the purpose of benefit and risk assessment (Baddock et al., 2019; Ball, 2006; McKenna & Gettler, 2016; Mobbs, Mobbs, & Mobbs, 2016).

As can be seen there have been a number of studies conducted over the past 20 years or more attempting to look at the relationships of maternal/caregiver sleep, hazardous risk factors, breastfeeding, and infant death. Many are limited by poor or inadequate data collection (e.g., unclear where the baby was positioned [such as bed or sofa], and in what position, at the time of death or during which part of the night);

that information regarding confounders like cigarette smoking, or drug and alcohol use was not collected; and lack of data regarding cultural and socioecological factors that differ between populations and must be taken into consideration.

There are many factors that we do know. We know the benefits of human milk and breastfeeding to infants and their mothers, and that these are dose-dependent; that breastfeeding through the night makes physiological sense because the largest prolactin surge in a 24 hr period occurs in the middle of the night; that breastfeeding through the night is associated with higher rates of exclusive breastfeeding and the longer duration of breastfeeding, which leads to higher doses of human milk. So, it is not a huge leap to accept that safe breastfeeding through the night that does not lead to an increase in infant death, hence "breastsleeping", is desirable. With current data, correctly interpreted, we are able to say this and develop public policy, in line with that already in place in the United Kingdom, to support families to safely sleep with and breastfeed their infants through the night. This concept is a potential game-changer given the current polarized debate on

Table 2. National Institute for Health and Care Excellence Guidance on Co-sleeping and Sudden Infant Death Syndrome Issued for England and Wales (2014).**Co-sleeping and Sudden Infant Death Syndrome**

The cause of sudden infant death syndrome (SIDS) is not known. It is possible that many factors contribute, but some factors are known to make SIDS more likely. These include placing a baby on her or his front or side to sleep. We need clear evidence to say that a factor directly causes SIDS. Evidence was reviewed relating to co-sleeping (parents or carers sleeping on a bed or sofa or chair with an infant) in the 1st year of an infant's life. Some of the reviewed evidence showed that there is a statistical relationship between SIDS and co-sleeping. This means that where co-sleeping occurs, there may be an increase in the number of cases of SIDS. However, the evidence does not allow us to say that co-sleeping causes SIDS. Therefore, the term association has been used in the recommendations to describe the relationship between co-sleeping and SIDS. The recommendations on co-sleeping and SIDS cover the 1st year of an infant's life.

1.4.47 Recognize that co-sleeping can be intentional or unintentional. Discuss this with parents and carers and inform them that there is an association between co-sleeping (parents or carers sleeping on a bed or sofa or chair with an infant) and SIDS.

1.4.48 Inform parents and carers that the association between co-sleeping (sleeping on a bed or sofa or chair with an infant) and SIDS is likely to be greater when they, or their partner, smoke.

1.4.49 Inform parents and carers that the association between co-sleeping (sleeping on a bed or sofa or chair with an infant) and SIDS may be greater with

- parental or carer recent alcohol consumption, or
- parental or carer drug use, or
- low birth weight or premature infants.

Note. Adapted from "Guidance on Cosleeping and Sudden Infant Death Syndrome Issued for England and Wales," by the National Institute for Health and Care Excellence, 2014. Copyright 2019 by the National Institute for Health and Care Excellence.

what we should be advising parents. In many places health-care providers, including physicians, nurses, and International Board Certified Lactation Consultants, are hampered by current policy in that they are not able to counsel families on safe sleep practices. This evokes fear of sleeping on the safe surface of their bed and puts families at risk of getting out of bed to feed their baby on a chair or sofa, and falling asleep there instead—a significantly more unsafe place. The onus is on us to call for better designed studies, with adequate interpretation, and appropriate translation into public policy for the safety of families globally.

Risk-Reduction Strategies

Different strategies have been adopted to advise parents on bed sharing over the last decade. One (Table 1) is to advise against bed sharing which has been adopted in some countries, such as the United States since 2005 (American Academy of Pediatrics, 2005; Moon, 2016a). Another strategy, adopted in some countries, for example the UK (Table 2), is to acknowledge that bed sharing occurs either intentionally or unintentionally and that it is therefore appropriate to discuss circumstances in which it is risky to bed-share (National Institute for Health and Care Excellence, 2014b).

The US strategy has the advantage of being a clear direct message to the public and therefore perceived to be an easier one to get across. Despite campaigns to decrease bed-sharing in some States, bed-sharing has increased in the US in recent years, especially among black and Hispanic communities (Colson et al., 2014). Taking into account the potential diagnostic shift currently happening in the US from SIDS to Accidental Suffocation and Strangulation in Bed (ASSB), the combined sudden unexpected infant death rate (SUID) appears to be almost flat-lining (Figure 2; CDC/NHCHS,

2019). There is now recognition that current AAP (2016) recommendations about bed-sharing are not being followed as widely as hoped, due to the complexity of the practice and the reasons for choosing to do it. Bed-sharing is a culturally ingrained infant care practice and in some low income communities, is used to keep infants safe (Joyner, Oden, Ajao, & Moon, 2010), while others choose to bed-share to facilitate breastfeeding. During one trial using enhanced messaging with high-risk families to avoid bed-sharing, the prevalence of bed-sharing actually increased rather than decreased (Moon et al., 2017).

Although the UK approach lacks the same simplicity as the US approach, it is more closely aligned to the evidence; acknowledging that bed-sharing happens means it can be discussed without judgement and specific hazardous situations or environments can be discussed (UNICEF UK, 2019). In the UK, both the SIDS rate and combined SUDI rate have fallen over the last 10 years (Figure 3; Office for National Statistics Great Britain, 2017), although it is not clear which elements of the risk reduction strategy have had the greater influence. The approach in the US in its most recent 2016 set of guidelines has been more nuanced; although they do not recommend bed-sharing they do acknowledge that mothers often fall asleep while breastfeeding their babies in bed, and advise that the parental bed should be prepared to avoid hazardous bedding should this occur (Moon, 2016a). The latest guidelines also advise that it is safer to breastfeed in bed at nighttime, compared to on sofas or armchairs, and strongly advise against the latter.

In New Zealand, the high SIDS rate among the Māori population, who often bed-share, and among whom smoking rates are high, has led to an intervention based on the premise that the provision of a separate sleep surface deployed in a shared sleep environment would reduce the risk of hazardous




Figure 2. Trends in US Sudden Unexpected Infant Death by cause, 1990–2017.




Figure 3. Trends in SIDS and SUDI in England and Wales (2000–2015).

bed-sharing. The *wahakura*, a woven flax bassinet-like structure and the *Pēpi-Pod™*, a plastic box of similar proportions,

were introduced into SIDS/SUDI prevention efforts in 2006 and 2011, respectively (Abel & Tipene-Leach, 2013). The

intervention involved the distribution of these devices in high SUDI risk situations, like smoking during pregnancy, and the provision of one-on-one counseling that heightened awareness of risk and promoted “Safe Sleep.” Importantly, the acceptability of this change to a culturally valued practice has been enhanced by the appeal of “reclaiming” a traditional Māori infant care practice (Abel, Stockdale-Frost, Rolls, & Tipene-Leach, 2015) and the fact that this device, placed in the shared bed, maintains the highly valued proximity of mother and infant. The high SIDS and SUDI rates in New Zealand have fallen dramatically since 2009, especially among Māori infants and in the areas with the most intensive Safe Sleep programs (Mitchell, Cowan, & Tipene-Leach 2016). Based on these results, the New Zealand Ministry of Health has recently adopted the Safe Sleep program as national policy.

The Way Forward

It is now clear that bed-sharing on its own does not substantially increase the risk of SIDS, but bed-sharing in conjunction with other hazardous circumstances—for instance smoking, alcohol consumption, drug use, and ad-hoc sleeping arrangements (e.g., sofa sharing) are clearly implicated (Blair, Sidebotham, Pease, & Fleming, 2014), within a larger context of poverty and inequality (Bartick & Tomori, 2018). Different risk reduction strategies have been used in different populations, although there are signs that a mixed-strategy approach targeting certain populations may be more beneficial in the future. What is clear is that the negative rhetoric that eliminates any hope of honest, bi-directional conversations between bed-sharing parents and their health providers must cease and be replaced by an emphasis on the magnitude of risk surrounding unsafe sleeping practices involving alcohol, drugs, and sofas or chairs, and establish a more coordinated approach with other public health strategists on how best to care for the infants as well as to keep them safe.

Declaration of Conflicting Interests

The authors declare the following potential conflicts of interest with respect to the research, authorship, and/or publication of this article: Dr Helen Ball served on the 2014 NICE Guidance Update Committee on Co-Sleeping & SIDS. She is Chair of the Lullaby Trust Scientific Committee and a Scientific Advisor to the Trust, and she has collaborated with both UNICEF UK Baby Friendly Initiative and Lullaby Trust in the production of safer sleep guidance for parents and health professionals in the UK. She has also served on the Board of ISPID (International Society for the Study and Prevention of Infant Deaths) and serves on the *JHL* Editorial Review Board. Professor Peter Blair is a Scientific Advisor to UNICEF United Kingdom Baby Friendly Initiative and helped write the safer sleep guidance for parents and health professionals in the United Kingdom. He is also chair of International Society for the Study and Prevention of Perinatal and Infant Deaths (ISPID). Dr Kathleen Marinelli serves as an Associate Editor for *JHL*. Professor James McKenna is an Emeritus Professor of Anthropology and the

Director of the Mother-Baby Behavioral Sleep Laboratory at the University of Notre Dame.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Kathleen A. Marinelli  <https://orcid.org/0000-0001-9342-6833>

References

- Abel, S., & Tipene-Leach, D. (2013). SUDI prevention: A review of Māori safe sleep innovations for infants. *New Zealand Medical Journal*, *126*(1379), 86–94.
- Abel, S., Stockdale-Frost, A., Rolls, R., & Tipene-Leach, D. (2015). The *wahakura*: A qualitative study of the flax bassinet as a sleep location for New Zealand Māori infants. *New Zealand Medical Journal*, *128*(1413), 12–19.
- American Academy of Pediatrics. (2005). The changing concept of sudden infant death syndrome: Diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in reducing risk. *Pediatrics*, *116*, 1245–55.
- American Academy of Pediatrics. (2019). Reducing sudden infant death with “Back to Sleep”. Retrieved from <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/7-great-achievements/Pages/Reducing-Sudden-Infant-Death-with-Back-to-.aspx>
- Ateah, C. A., & Hamelin, K. J. (2008). Maternal bedsharing practices, experiences, and awareness of risks. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, *37*(3), 274–281. doi:10.1111/j.1552-6909.2008.00242.x
- Baddock, S. A., Purnell, M. T., Blair, P. S., Pease, A. S., Elder, D. E., & Galland, B. C. (2019). The influence of bed-sharing on infant physiology, breastfeeding and behaviour: A systematic review. *Sleep Medicine Reviews*, *43*, 106–117.
- Ball, H. L. (2002). Reasons to bed-share: Why parents sleep with their infants. *Journal of Reproductive and Infant Psychology*, *20*(4), 207–221. doi:10.1080/0264683021000033147
- Ball, H. L. (2003). Breastfeeding, bed-sharing, and infant sleep. *Birth*, *30*(3), 181–188. doi:10.1046/j.1523-536X.2003.00243.x
- Ball, H. L. (2006). Parent-infant bed-sharing behavior. *Human Nature*, *17*(3), 301–318. doi:10.1007/s12110-006-1011-1
- Ball, H. L. (2007). Bed-sharing practices of initially breastfed infants in the first 6 months of life. *Infant and Child Development*, *16*, 387–401. doi:10.1002/icd.519
- Ball, H. L. (2008). Evolutionary paediatrics: A case study in applying Darwinian medicine. In S. Elton, & P. O’Higgins (Eds.), *Medicine and evolution: Current applications, future prospects*, pp. 125–50. London: Taylor & Francis.
- Ball, H. L. (2017a). The Atlantic divide: Contrasting UK and US recommendations on cosleeping and bed-sharing. *Journal of Human Lactation* *33*(4), 765–769. doi: 10.1177/0890334417713943.
- Ball, H. L. (2017b). Evolution-informed maternal-infant health. *Nature Ecology and Evolution*, *1*(3), 1–3. doi:10.1038/s41559-017-0073
- Ball, H. L., Hooker, E., & Kelly, P. J. (1999). Where will the baby sleep? Attitudes and practices of new and experienced

- parents regarding cosleeping with their newborn infants. *American Anthropologist*, 101(1), 143–151. doi:10.1525/aa.1999.101.1.143
- Ball, H. L., Howel, D., Bryant, A., Best, E., Russell, C., & Ward-Platt, M. (2016). Bed-sharing by breastfeeding mothers: Who bed-shares and what is the relationship with breastfeeding duration? *Acta Paediatrica*, 105(6), 628–634. doi:10.1111/apa.13354
- Ball, H. L., Moya, E., Fairley, L., Westman, J., Oddie, S., & Wright, J. (2012). Infant care practices related to sudden infant death syndrome in South Asian and white British families in the UK. *Paediatric and Perinatal Epidemiology*, 26(1), 3–12. doi:10.1111/j.1365-3016.2011.01217.x
- Ball, H. L., & Russell, C. K. (2012). Nighttime nurturing: An evolutionary perspective on breastfeeding and sleep. In D. Narvaez, J. Panksepp, A. Schore, & T. Gleason (Eds.), *Evolution, early experience and human development: From research to practice and policy* (pp. 241–261). Oxford University Press.
- Ball, H. L., & Volpe, L. E. (2013). Sudden Infant Death Syndrome (SIDS) risk reduction and infant sleep location: Moving the discussion forward. *Social Science and Medicine*, 79, 84–91. doi:10.1016/j.socscimed.2012.03.025
- Ball, H. L., Ward-Platt, M. P., Heslop, E., Leech, S. J., & Brown, K. A. (2006). Randomised trial of infant sleep location on the postnatal ward. *Archives of Diseases in Childhood* 91(12), 1005–10.
- Bartick, M., & Smith, L. J. (2014). Speaking out on safe sleep: Evidence-based infant sleep recommendations. *Breastfeeding Medicine*, 9(9), 417–22. doi:10.1089/bfm.2014.0113
- Bartick, M., & Tomori, C. (2018). Sudden infant death and social justice: A syndemics approach. *Maternal & Child Nutrition*, e12652. doi:10.1111/mcn.12652
- Blair, P. S., & Ball, H. L. (2004). The prevalence and characteristics associated with parent-infant bed-sharing in England. *Archives of Diseases in Childhood*, 89(12), 1106–10. doi:10.1136/ad.2003.038067
- Blair, P. S., Heron, J., & Fleming, P. J. (2010). Relationship between bed sharing and breastfeeding: Longitudinal, population-based analysis. *Pediatrics*, 126(5), e1119–e1126. doi:10.1542/peds.2010-1277
- Blair, P. S., Sidebotham, P., Berry, P. J., Evans, M., & Fleming, P. J. (2006). Major epidemiological changes in sudden infant death syndrome: A 20-year population-based study in the UK. *Lancet* 367, (9507) 314–9.
- Blair, P. S., Sidebotham, P., Evason-Coombe, C., Edmonds, M., Heckstall-Smith, E. M., & Fleming, P. (2009). Hazardous cosleeping environments and risk factors amenable to change: Case-control study of SIDS in south west England. *British Medical Journal*, 339, b3666.
- Blair, P. S., Sidebotham, P., Pease, A., & Fleming, P. J. (2014). Bed-sharing in the absence of hazardous circumstances: Is there a risk of Sudden Infant Death Syndrome? An analysis from two case-control studies conducted in the UK. *PloS One*, 9(9), e107799. doi:10.1371/journal.pone.0107799
- Carpenter, R., McGarvey, C., Mitchell, E. A., Tappin, D. M., Vennemann, M. M., Smuk, M., & Carpenter, J. R. (2013). Bed sharing when parents do not smoke: Is there a risk of SIDS? An individual level analysis of five major case-control studies. *British Medical Journal Open*, 3(5), e002299.
- Center for Disease Control and Prevention/National Center for Health Statistics, Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion National Vital Statistics System, & (CDC/NHCHS) Compressed Mortality File. (2019). *Trends in US Sudden Unexpected Infant Death by Cause, 1990-2017*. Retrieved from <https://www.cdc.gov/sids/data.htm>
- Colson, E. R., Willinger, M., Rybin, D., Heeren, T., Smith, L. A., Lister, G., & Corwin, M. J. (2013). Trends and factors associated with infant bed sharing, 1993–2010. *Journal of the American Medical Association Pediatrics*, 167(11), 1032. doi:10.1001/jamapediatrics.2013.2560
- Côté, A. (2006). Bed sharing and sudden infant death syndrome: Do we have a definition problem? *Paediatrics and Child Health*, 11, 34–38.
- Crane, D., & Ball, H. L. (2016). A qualitative study in parental perceptions and understanding of SIDS-reduction guidance in a UK bi-cultural urban community. *BMC Pediatrics*, 16(1), 23. doi:10.1186/s12887-016-0560-7
- Culver, E. D. (2009). Exploring bed-sharing mothers' motives and decision-making for getting through the night intact: A grounded theory. *Journal of Midwifery and Women's Health*, 54(5), 423–423. doi:10.1016/j.jmwh.2009.06.010
- Fetherston, C. M., & Leach, J. S. (2012). Analysis of the ethical issues in the breastfeeding and bedsharing debate. *Breastfeeding Review*, 20(3), 7–17.
- Gilbert, R., Salanti, G., Harden, M., & See, S. (2005). Infant sleeping position and the sudden infant death syndrome: Systematic review of observational studies and historical review of recommendations from 1940 to 2002. *International Journal of Epidemiology*, 34(4), 874–87. doi:10.1093/ije/dyi088
- Hauck, F. R., Thompson, J. M. D., Tanabe, K. O., Moon, R. Y., & Vennemann, M. M. (2011). Breastfeeding and reduced risk of sudden infant death syndrome: A meta-analysis. *Pediatrics*, 128(1), 103–110. doi:10.1542/peds.2010-3000
- Jelliffe, D. B., & Jelliffe, E. F. P. (1978). Human milk in the modern world. *British Medical Journal*, 2(6151), 1573.
- Joyner, B. L., Oden, R. P., Ajao, T. I., & Moon, R. Y. (2010). Where should my baby sleep: A qualitative study of African American infant sleep location decisions. *Journal of the National Medical Association*, 102(10), 81–9.
- Konner, M., & Super, C. (1987). Sudden Infant Death Syndrome: An anthropological hypothesis. In C. M. Super (Ed.), *The role of culture in developmental disorder*, pp. 95–108. San Diego: Academic Press.
- Luijk, M. P. C.M., Mileva-Seitz, V. R., Jansen, P. W., van IJzendoorn, M. H., Jaddoe, V. W. V., Raat, H., & ...Tiemeier, H. (2013). Ethnic differences in prevalence and determinants of mother-child bed-sharing in early childhood. *Sleep Medicine*, 14(11), 1092–9. doi:10.1016/j.sleep.2013.04.019
- McCoy, R. C., Hunt, C. E., Lesko, S. M., Vezina, R., Corwin, M. J., Willinger, M., ...Mitchell, A. A. (2004). Frequency of bed sharing and its relationship to breastfeeding. *Journal of Developmental and Behavioral Pediatrics*, 25(3), 141–9.
- McKenna, J. J. (1986). An anthropological perspective on the sudden infant death syndrome (SIDS): The role of parental breathing cues and speech breathing adaptations. *Medical Anthropology*, 10(1), 9–92. doi:10.1080/01459740.1986.9965947
- McKenna, J. J., Ball, H. L., & Gettler, L. T. (2007). Mother-infant cosleeping, breastfeeding and Sudden Infant Death Syndrome: What biological anthropology has discovered about normal infant sleep and pediatric sleep medicine. *American Journal of Physical Anthropology*, 134(S45), 133–161. doi:10.1002/ajpa.20736

- McKenna, J. J., & Gettler, L. T. (2016). There is no such thing as infant sleep, there is no such thing as breastfeeding, there is only breastsleeping. *Acta Paediatrica*, *105*(1):17–21. doi:10.1111/apa.13161.
- McKenna, J. J., Mosko, S., Dungy, C., & McAninch, J. (1990). Sleep and arousal patterns of co-sleeping human mother/infant pairs: A preliminary physiological study with implications for the study of sudden infant death syndrome (SIDS). *American Journal of Physical Anthropology*, *83*(3), 331–47. doi:10.1002/ajpa.1330830307
- McKenna, J. J., Thoman, E. B., Anders, T. F., Sadeh, A., Schechtman, V. L., & Glotzbach, S. F. (1993). Infant-parent co-sleeping in an evolutionary perspective: Implications for understanding infant sleep development and the sudden infant death syndrome. *Sleep*, *16*(3), 263–82.
- McKenna, J. J., & Volpe, L. (2007). An internet based study of infant sleeping arrangements and parental perceptions. *Infant and Child Development* *16*, 359–385.
- Mitchell, E. A., Cowan, S., & Tipene-Leach, D. (2016). The recent fall in postperinatal mortality in New Zealand and the Safe Sleep programme. *Acta Paediatrica*, *105*(11), 312–320.
- Mitchell, E. A., & Scragg, R. (1993). Are infants sharing a bed with another person at increased risk of sudden infant death syndrome? *Sleep*, *16*(4), 387–9.
- Mitchell, E., Thompson, J., Zuccollo, J., MacFarlane, M., Taylor, B., Elder, D., ... Fleming, P. (2017). The combination of bed-sharing and maternal smoking leads to a greatly increased risk of sudden unexpected death in infancy: The New Zealand SUDI nationwide case control study. *New Zealand Medical Journal*, *130*(1456), 52–64.
- Mobbs, E., Mobbs, G., & Mobbs, A. (2016). Imprinting, latchment and displacement: A review of early instinctual behaviour influencing breastfeeding success. *Acta Paediatrica*, *105*, 24–30.
- Moon, R. Y., Mathews, A., Joyner, B. L., Oden, R. P., He, J., & McCarter, R. (2017). Health messaging and African-American infant sleep location: A randomized controlled trial. *Journal of Community Health*, *42*(1), 1–9.
- Moon, R. Y., & The Task Force on Sudden Infant Death Syndrome. (2016a). SIDS and other sleep-related infant deaths: Updated 2016 recommendations for a safe infant sleeping environment. *Pediatrics*, *138*(5), e20162938.
- Moon, R. Y. & The Task Force on Sudden Infant Death Syndrome. (2016b). SIDS and other sleep related infant deaths: Evidence base for the 2016 updated recommendations for a safe infant sleeping environment. *Pediatrics*. *138*(5), e20162940.
- Mosko, S., McKenna, J., Dickel, M., & Hunt, L. (1993). Infant-parent cosleeping: The appropriate context for the study of infant sleep and implications for sudden infant death syndrome (SIDS) research. *Journal of Behavioral Medicine*, *16*(6), 589–610.
- Mosko, S., Richard, C., & McKenna, J. (1997a). Infant arousals during mother-infant bed sharing: Implications for infant sleep and sudden infant death syndrome research. *Pediatrics*, *100*(5), 841–9.
- Mosko, S., Richard, C., & McKenna, J. (1997b) Maternal sleep and arousals during bedsharing with infants. *Sleep*, *20*(2), 142–150.
- National Institute for Health and Care Excellence. (2014a). Addendum to clinical guideline 37, postnatal care. Retrieved from <https://www.nice.org.uk/guidance/cg37/evidence/full-guideline-addendum-485782238>
- National Institute for Health and Care Excellence. (2014b). Postnatal care tools and resources. Retrieved from <https://www.nice.org.uk/guidance/cg37/resources>
- Office for National Statistics Great Britain. (2017). Data retrieved by private request from <https://www.ons.gov.uk/>
- Quillin, S. I. M., & Glenn, L. L. (2004). Interaction between feeding method and co-sleeping on maternal-newborn sleep. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, *33*(5), 580–8.
- Rechtman, L. R., Colvin, J. D., Blair, P. S., & Moon, R. Y. (2014). Sofas and infant mortality. *Pediatrics*, *134*(5), e1293–300.
- Rudzik, A. E. F., & Ball, H. L. (2016). Exploring maternal perceptions of infant sleep and feeding method among mothers in the United Kingdom: A qualitative focus group study. *Maternal and Child Health Journal*, *20*(1), 33–40. doi:10.1007/s10995-015-1798-7
- Salm Ward, T. C. (2015). Reasons for mother-infant bed-sharing: A systematic narrative synthesis of the literature and implications for future research. *Maternal and Child Health Journal*, *19*(3), 675–90. doi:10.1007/s10995-014-1557-1
- Salm Ward, T. C., & Doering, J. J. (2014). Application of a socio-ecological model to mother-infant bed-sharing. *Health Education and Behavior*, *41*(6), 577–589. doi:10.1177/1090198114543010
- Santos, I. S., Mota, D. M., Matijasevich, A., Barros, A. J. D., & Barros, F. C. F. (2009). Bed-sharing at 3 months and breastfeeding at 1 year in Southern Brazil. *Journal of Pediatrics*, *155*(4), 505–509. doi:10.1016/j.jpeds.2009.04.037
- Scragg, R., Mitchell, E. A., Taylor, B. J., Stewart, A. W., Ford, R. P., Thompson, J. M., Allen, E. M., & Becroft, D.M. (1993). Bed sharing, smoking, and alcohol in the sudden infant death syndrome. New Zealand Cot Death Study Group. *British Medical Journal*, *307*(6915), 1312–8.
- Scragg, R. K. R., Stewart, A. W., Mitchell, E. A., Thompson, J. M. D., Taylor, B. J. B. J., Williams, S. M., & ...Thompson, J. M. D. (1996). Infant room-sharing and prone sleep position in sudden infant death syndrome. *The Lancet*, *347*(8993), 7–12. doi:10.1016/S0140-6736(96)91554-8
- Smith, L. A., Geller, N. L., Kellams, A. L., Colson, E. R., Rybin, D. V., Heeren, T., & Corwin, M. J. (2016). Infant sleep location and breastfeeding practices in the United State: 2011–2014. *Academic Pediatrics*. *16*, 540–549.
- Thompson, J. M. D., Tanabe, K., Moon, R. Y., Mitchell, E. A., McGarvey, C., Tappin, D., ...Hauck, F. R. (2017). Duration of breastfeeding and risk of SIDS: An individual participant data meta-analysis. *Pediatrics*, *140*(5), e20171324. doi:10.1542/peds.2017-1324
- Tully, K. P., & Ball, H. L. (2013). Trade-offs underlying maternal breastfeeding decisions: A conceptual model. *Maternal and Child Nutrition*, *9*(1), 90–98. doi:10.1111/j.1740-8709.2011.00378.x
- UNICEF UK, Baby Friendly Initiative. (2019). *Caring for your baby at night, a guideline for parents*. Retrieved from <https://www.unicef.org.uk/babyfriendly/wp-content/uploads/sites/2/2018/08/Caring-for-your-baby-at-night-web.pdf>
- Vennemann, M. M., Hense, H. W., Bajanowski, T., Blair, P. S., Complojer, C., Moon, R. Y., & Kiechl-Kohlendorfer, U. (2012). Bed sharing and the risk of Sudden Infant Death Syndrome: Can we resolve the debate? *Journal of Pediatrics*, *160*(1), 44–8.
- Willinger, M., Ko, C., & Hoffman, H. (2003). Trends in infant bed sharing in the United States, 1993–2000: The National Infant Sleep Position study. *Archives of Pediatrics*, *157*, 43–49.